

Math 17B

Vogler

### Estimating the Value of a Definite Integral

Suppose that the integral  $\int_a^b f(x) dx$  is too difficult (or impossible) to compute, or that you are simply required to estimate its exact value. The following two methods offer two different ways to determine good estimates.

#### 1.) MIDPOINT RULE

- Divide the interval  $[a, b]$  into  $n$  equal parts, each of length  $h = \frac{b-a}{n}$ .
- Let  $a = x_0, x_1, x_2, x_3, \dots, x_{n-1}, x_n = b$  be the partition of the interval and let the sampling points  $c_1, c_2, c_3, \dots, c_n$  be the MIDPOINTS of these subintervals.

c.) The Midpoint Estimate for  $\int_a^b f(x) dx$  is

$$M_n = h [f(c_1) + f(c_2) + f(c_3) + \dots + f(c_n)] .$$

d.) The Absolute Error  $|E_n| = \left| \int_a^b f(x) dx - M_n \right|$  satisfies

$$|E_n| \leq (b-a) \frac{h^2}{24} \left\{ \max_{a \leq x \leq b} |f''(x)| \right\} = \frac{(b-a)^3}{24n^2} \left\{ \max_{a \leq x \leq b} |f''(x)| \right\} .$$

#### 2.) TRAPEZOIDAL RULE

- Divide the interval  $[a, b]$  into  $n$  equal parts, each of length  $h = \frac{b-a}{n}$ .
- Let  $a = x_0, x_1, x_2, x_3, \dots, x_{n-1}, x_n = b$  be the partition of the interval.
- The Trapezoidal Estimate for  $\int_a^b f(x) dx$  is

$$T_n = \frac{h}{2} [f(x_0) + 2f(x_1) + 2f(x_2) + \dots + 2f(x_{n-1}) + f(x_n)] .$$

d.) The Absolute Error  $|E_n| = \left| \int_a^b f(x) dx - T_n \right|$  satisfies

$$|E_n| \leq (b-a) \frac{h^2}{12} \left\{ \max_{a \leq x \leq b} |f''(x)| \right\} = \frac{(b-a)^3}{12n^2} \left\{ \max_{a \leq x \leq b} |f''(x)| \right\} .$$

Ex Determine the value of  $n$  so that the Trapezoid Estimate  $T_n$  estimates the exact value of  $\int_{-1}^2 \frac{1}{\sqrt{x+5}} dx$  with absolute error at most 0.0001.

Let  $f(x) = \frac{1}{\sqrt{x+5}} = (x+5)^{-\frac{1}{2}}$  on  $[-1, 2]$ . Then  
 $h = \frac{b-a}{n} = \frac{2-(-1)}{n} = \frac{3}{n}$  and  $f'(x) = -\frac{1}{2}(x+5)^{-\frac{3}{2}}$   
 $\Rightarrow f''(x) = \frac{3}{4}(x+5)^{-\frac{5}{2}} = \frac{3}{4(x+5)^{\frac{5}{2}}}.$

We have  $\max_{-1 \leq x \leq 2} |f''(x)| = \max_{-1 \leq x \leq 2} \left| \frac{3}{4(x+5)^{\frac{5}{2}}} \right|$   
 $= \max_{-1 \leq x \leq 2} \frac{3}{4|x+5|^{\frac{5}{2}}} = \frac{3}{4|-1+5|^{\frac{5}{2}}} = \frac{3}{128}.$

Hence,  $|E_n| \leq \frac{(b-a)^3}{12n^2} \left\{ \max_{-1 \leq x \leq 2} |f''(x)| \right\} = \frac{3^3}{12n^2} \left\{ \frac{3}{128} \right\}$

$$= \frac{27}{512} \cdot \frac{1}{n^2} \leq 0.0001$$

$$\Rightarrow n^2 \geq \frac{27}{512(0.0001)} \Rightarrow n \geq \sqrt{\frac{27}{512(0.0001)}} \approx 22.9$$

so choose  $\boxed{n=23}$  (or bigger integer).