

Math 17B
Vogler
Taylor Polynomials

QUESTION : Consider the function $y = f(x)$ and assume that mathematical circumstances would require that this function be replaced with an n th-degree polynomial whose values closely approximate the values of $y = f(x)$. How would we determine the coefficients of this unknown polynomial ?

ANSWER : Assume that $y = f(x)$ is a given function and constant “ a ” is known. Determine a list of real numbers $a_0, a_1, a_2, a_3, \dots, a_n$ so that the polynomial is

$$P_n(x) = a_0 + a_1(x - a) + a_2(x - a)^2 + a_3(x - a)^3 + \dots + a_n(x - a)^n ,$$

which satisfies the condition

$$f(x) \approx P_n(x) ,$$

i.e.,

$$(T) \quad f(x) \approx a_0 + a_1(x - a) + a_2(x - a)^2 + a_3(x - a)^3 + \dots + a_n(x - a)^n .$$

If we assume that “ \approx ” is “ $=$ ” and substitute $x = a$ in equation (T), we get

$$f(a) = a_0 + a_1(0) + a_2(0)^2 + a_3(0)^3 + \dots + a_n(0)^n = a_0 ,$$

i.e.,

$$a_0 = f(a) .$$

Now differentiate equation (T) term by term getting

$$f'(x) = a_1 + 2a_2(x - a) + 3a_3(x - a)^2 + 4a_4(x - a)^3 + \dots + na_n(x - a)^{n-1} .$$

If we substitute $x = a$ in this equation, we get

$$f'(a) = a_1 + 2a_2(0) + 3a_3(0)^2 + 4a_4(0)^3 + \dots + na_n(0)^{n-1} = a_1 ,$$

i.e.,

$$a_1 = f'(a) .$$

Now differentiate again term by term getting

$$f''(x) = 2a_2 + 3 \cdot 2a_3(x - a) + 4 \cdot 3a_4(x - a)^2 + 5 \cdot 4a_5(x - a)^3 + \dots + n \cdot (n - 1)a_n x^{n-2} .$$

If we substitute $x = a$ in this equation, we get

$$f''(a) = 2a_2 + 3 \cdot 2a_3(0) + 4 \cdot 3a_4(0)^2 + 5 \cdot 4a_5(0)^3 + \dots + (n - 1) \cdot na_n(0)^{n-2} = 2a_2 ,$$

i.e.,

$$a_2 = \frac{f''(a)}{2!} .$$

Now differentiate again term by term getting

$$f'''(x) = 3 \cdot 2a_3 + 4 \cdot 3 \cdot 2a_4(x - a) + 5 \cdot 4 \cdot 3a_5(x - a)^2 + \dots + n(n - 1)(n - 2)a_n x^{n-3} .$$

If we substitute $x = a$ in this equation, we get

$$f'''(a) = 3 \cdot 2a_3 + 4 \cdot 3 \cdot 2a_4(0) + 5 \cdot 4 \cdot 3a_5(0)^2 + \cdots n(n-1)(n-2)a_n(0)^{n-3} = 3 \cdot 2a_3 ,$$

i.e.,
$$a_3 = \frac{f'''(a)}{3!} .$$

Continuing this term by term differentiation and substitution process results in the fact that

$$(S) \quad a_k = \frac{f^{(k)}(a)}{k!} \quad \text{for } k = 0, 1, 2, 3, \dots, n .$$

DEFINITION : The *Taylor Polynomial of degree n* centered at $x = a$ for function $y = f(x)$ is given by

$$(P) \quad P_n(x) = a_0 + a_1(x - a) + a_2(x - a)^2 + a_3(x - a)^3 + \cdots + a_n(x - a)^n$$

and

$$(S) \quad a_k = \frac{f^{(k)}(a)}{k!} \quad \text{for } k = 0, 1, 2, 3, \dots, n .$$

QUESTION : Consider an ordinary function $y = f(x)$ and $P_n(x)$, its Taylor Polynomial of degree n centered at $x = a$. We assume that $f(x) \approx P_n(x)$. Let the error be given by $R_{n+1}(x; a) = f(x) - P_n(x)$. It can be shown that the absolute error (Taylor Error) for their difference is

$$\left| R_{n+1}(x; a) \right| = \left| \frac{f^{(n+1)}(c)}{(n+1)!} (x - a)^{n+1} \right| ,$$

where c is some number between x and a . (Note that the number c appears as a result of using the Intermediate Value Theorem to derive this error formula.)